

Global Fits of the MSSM and SM to Electroweak Precision Data

W. DE BOER

*Inst. für Experimentelle Kernphysik, Univ. of Karlsruhe,
Postfach 6980, 76128 Karlsruhe, Germany*

The Minimal supersymmetric extension of the Standard Model (MSSM) with light stops, charginos or pseudoscalar Higgs bosons has been suggested as an explanation of the too high value of the branching ratio of the Z^0 boson into b quarks (R_b anomaly). A program including all radiative corrections to the MSSM at the same level as the radiative corrections to the SM has been developed and used to perform global fits to all electroweak data from LEP, SLC and the Tevatron. Recent updates of electroweak data, presented at this conference, reduce the R_b anomaly from a 3.2σ to a 1.8σ effect. In addition, the $b \rightarrow s\gamma$ decay is 30% below the SM prediction. In the MSSM light stops and light charginos increase R_b and decrease the $b \rightarrow s\gamma$ rate, so both observations can be brought into agreement with the MSSM for the same region of parameter space. However, the resulting χ^2 value for the MSSM fits is only marginally lower. In addition, the splitting in the stop sector has to be unnaturally high, so it remains to be seen if these effects are real or due to a fluctuation.

1 Introduction

Previous LEP data showed a too high value of R_b (3.2σ) and a too low value of R_c (2σ) where $R_{b(c)}$ is the ratio $R_{b(c)} = \Gamma_{Z^0 \rightarrow b\bar{b}(c\bar{c})} / \Gamma_{Z^0 \rightarrow q\bar{q}}$. In the past it has been shown by several groups that it is possible to increase R_b using minimal supersymmetric models (MSSM) with light charginos, stops or Higgses, which yield positive contributions to the $Zb\bar{b}$ vertex, although no improvement in R_c could be obtained^{1,9}. Recent updates of electroweak data¹⁰ show no significant deviation of R_c from the Standard Model (SM) prediction, and a value of R_b which is 1.8σ above the SM value. Such a moderate deviation in R_b can either be a fluctuation or may originate from genuine MSSM contributions. In this paper an equivalent analysis of all electroweak data, both in the SM and its supersymmetric extension, is described using all actual electroweak data from Tevatron, LEP and SLC¹⁰, the measurement of $\frac{BR(b \rightarrow s\gamma)}{BR(b \rightarrow ce\bar{\nu})}$ from CLEO¹¹ and limits on the masses of supersymmetric particles. Further details of the procedure and extensive references are given elsewhere¹².

2 Results

2.1 Standard Model Fits

The SM cross sections are determined by $M_Z, m_t, m_h, G_F, \alpha, \alpha_s$. From the combined CDF and D0 data m_t has been determined to be 175 ± 6 GeV¹⁰, so the parameters with the largest uncertainties are m_h and α_s . The error on the finestruc-

ture constant α is limited by the uncertainty in the hadronic cross section in e^+e^- annihilation at low energies, which is used to determine the vacuum polarization contributions to α . The error was taken into account by considering α to be a free parameter in the fit and constraining it to the value $1/\alpha = 128.89 \pm 0.09$ ³. If this error is not taken into account, the error on the Higgs mass is underestimated by 30%. Using the input values discussed in the introduction yields:

$$\begin{aligned}\alpha_s &= 0.120 \pm 0.003 \\ m_t &= 172.0^{+5.8}_{-5.7} \text{ GeV} \\ m_H &= 141^{+140}_{-77} \text{ GeV} \\ \sin^2 \theta_{\overline{MS}} &= 0.2316 \pm 0.0004.\end{aligned}$$

Minor deviations from the EWWG fit results¹⁰ are due to the incorporation of the $b \rightarrow \gamma$ data from CLEO¹¹, which are important for the MSSM fits described below. The value of $\sin^2 \theta_{\overline{MS}}$ is within errors equal to $\sin^2 \Theta_{eff}^{lept}$. The main contributions to the $\chi^2/d.o.f = 19.6/15$ originate from $\sin^2 \Theta_{eff}^{lept}$ from SLD ($\Delta\chi^2 = 4.9$), R_b ($\Delta\chi^2 = 3.1$) and A_{FB}^b ($\Delta\chi^2 = 3.5$). The SM agreement is good for all other observables, including R_c , as shown in fig. 2.

The dependence of $\sin^2 \Theta_{eff}^{lept}$ on the SM Higgs mass is approximately logarithmic (see fig. 1). The LEP data alone without SLD yield $m_H \approx 240$ GeV, while $\sin^2 \Theta_{eff}^{lept}$ from SLD corresponds to $m_H \approx 15$ GeV, as indicated by the squares in fig. 1. The latter value is excluded by the lower limit of 58.4 GeV from the combined LEP exper-




Figure 1: Dependence of the SM $\sin^2 \Theta_{eff}^{lept}$ on the Higgs mass. The top mass $m_t = 175 \pm 6$ GeV was varied within its error, as shown by the dashed band labelled SM. The SLD and the LEP measurement of $\sin^2 \Theta_{eff}^{lept}$ are also shown as horizontal bands. The SLD value yields a Higgs mass below the recent limits by direct Higgs searches at LEP (shaded area).

iments¹⁴. The different values of $\sin^2 \Theta_{eff}^{lept}$ from LEP and SLD translate into different predictions for M_W , as shown in fig. 3. The present M_W measurements, including the preliminary value from the LEP II measurements at 161 GeV¹⁵, lie in between these predictions.

2.2 MSSM Fits and Comparison with the SM

As mentioned in the introduction, the MSSM can increase the value of R_b , which experimentally is slightly above the SM value. The major additional contributions originate from vertex contributions with light charginos and light right handed stops in the low $\tan \beta$ scenario and light higgses for large $\tan \beta$ values. Since the large $\tan \beta$ scenario does not improve R_b significantly¹², it will not be discussed here anymore. The R_b dependence on chargino and stop masses is shown in fig. 4. The experimental value $R_b = 0.2178 \pm 0.0011$ is clearly above the SM value of 0.2158 and can be obtained for charginos around 85 GeV and stop masses around 50 GeV (best fit results).

The fit results are compared with the Standard Model fits in fig. 2. The Standard Model




Figure 2: Fit results normalized to the SM- and MSSM ($\tan \beta = 1.6$) values. The difference in $\chi^2/d.o.f$ between the SM and MSSM originates mainly from R_b .




Figure 3: A compilation of the W-masses. The vertical lines indicate the predictions from the LEP I and SLD electroweak data determining $\sin^2 \Theta_{eff}^{lept}$ and the data points represent the various direct measurements of M_W . The preliminary LEP II value was taken from presented at the LEPC Meeting¹⁵.




Figure 4: R_b as function of the stop and chargino masses. The experimental value $R_b = 0.2178 \pm 0.0011$ is clearly above the SM value of 0.2158 and can be obtained for light charginos and stops.




Figure 5: Comparison of α_s determinations with the fit results. The data have been taken from ¹⁴ and ¹⁶. Note the remarkable agreement between the new precise determinations: the values from the SM and MSSM fits and the new values from deep inelastic scattering and $b\bar{b}$ lattice calculations.

$\chi^2/d.o.f. = 18.5/15$ corresponds to a probability of 24%, the MSSM $\chi^2/d.o.f. = 16.1/12$ to a probability of 19%. In counting the d.o.f the insensitive (and fixed) parameters were ignored.

Another interesting point are the $\alpha_s(M_Z)$ values. An increase in R_b implies an increase in the total width of the Z^0 boson, which can be compensated by a decrease in the QCD corrections, i.e. α_s . Fig. 5 shows a comparison of different measurements of $\alpha_s(M_Z)$ with the fitted values given in this paper. The agreement is remarkable. Note that the α_s crisis has disappeared after the LEP value from the total cross section came down and the value from both lattice calculations and deep inelastic scattering went up¹⁶.

3 CMSSM and R_b

An increase in R_b requires one (mainly right handed) stop to be light and the other one to be heavy. If both would be light, then all other squarks are likely to be light, which would upset the good agreement between the SM and all other electroweak data. A large mass splitting in the stop sector needs a very artificial fine tuning of the few free parameters in the Constrained MSSM, which assumes unification of gauge and b- τ Yukawa couplings¹⁷. This is obvious from the mixing matrix:

$$\begin{pmatrix} \tilde{m}_L^2 + m_t^2 + D_L & m_t(A_t m_0 - \mu \cot \beta) \\ m_t(A_t m_0 - \mu \cot \beta) & \tilde{m}_R^2 + m_t^2 + D_R \end{pmatrix} \quad (1)$$

The D-terms proportional to $\cos 2\beta$ are negligible for $\tan \beta \approx 1$. If one of the diagonal elements is much larger than m_t , the off-diagonal terms of the order m_t will not cause a mixing and the difference between the left- and right-handed stops has to come from the evolution of the diagonal terms:

$$\begin{aligned} \frac{d\tilde{m}_L^2}{dt} &= \left(\frac{16}{3}\tilde{\alpha}_3 M_3^2 + 3\tilde{\alpha}_2 M_2^2 + \frac{1}{15}\tilde{\alpha}_1 M_1^2 \right) \quad (2) \\ &\quad - [Y_t(\tilde{m}_{Q_3}^2 + \tilde{m}_{U_3}^2 + m_{H_2}^2 + A_t^2 m_0^2) \\ &\quad + Y_b(\tilde{m}_{Q_3}^2 + \tilde{m}_{D_3}^2 + m_{H_1}^2 + A_b^2 m_0^2)] \end{aligned}$$

$$\begin{aligned} \frac{d\tilde{m}_R^2}{dt} &= \left(\frac{16}{3}\tilde{\alpha}_3 M_3^2 + \frac{16}{15}\tilde{\alpha}_1 M_1^2 \right) \quad (3) \\ &\quad - 2Y_t(\tilde{m}_{Q_3}^2 + \tilde{m}_{U_3}^2 + m_{H_2}^2 + A_t^2 m_0^2) \end{aligned}$$

One observes that the difference between left- and right handed stops depends on the Yukawa couplings for top and bottom (Y_t, Y_b) and the trilinear couplings $A_{t(b)}$. For low $\tan \beta$ Y_b is negligible,

while A_t and Y_t are not free parameters, since they go to fixed point solutions¹⁷. Therefore there is no freedom to adjust these parameters within the CMSSM in order to get a large splitting between the left- and right-handed stops.

In addition, problems arise with electroweak symmetry breaking, since this requires the Higgs mixing parameter μ to be much heavier than the gaugino masses¹⁷, while R_b requires low values of μ for a significant enhancement (since the chargino has to be preferably Higgsino-like). In conclusion, within the CMSSM an enhancement of R_b above the SM is practically excluded.

4 Conclusions

Both the MSSM and SM provide a good description of all electroweak data. With all mass bounds and the $b \rightarrow s\gamma$ rate included in the fit, the best $\chi^2/d.o.f$ in the MSSM (SM) is 16.1/12 (18.5/15), which corresponds to a probability of 19% (24%). The slightly better χ^2 of the MSSM is mainly due to the better description of R_b , but this requires an unnatural large splitting in the stop sector. Final analysis of available LEP data will teach of the present preliminary value of R_b will indeed stay above the SM value.

5 Acknowledgments

Thanks go to the Electroweak Working Group and the LEP experiments, who made the preliminary data available before the Conference and to my close collaborators Wolfgang Hollik and Ulrich Schwickerath during this analysis. This work was done during a sabbatical and support from the Volkswagen-Stiftung (Contract I/71681) is greatly appreciated.

6 References

1. *Proceedings of the Workshop Physics at LEP2*, Editors G. Altarelli, T. Sjöstrand, F. Zwirner, Vol.1 and Vol.2, CERN 96-01
2. M. Boulware, D. Finnell, *Radiative Corrections to $BR(Z \rightarrow b\bar{b})$ in the Minimal Supersymmetric Standard Model*, *Phys. Rev. D***44**(1991) 2054
3. P. H. Chankowski, S. Pokorski, *Chargino Mass and R_b Anomaly*, IFT-96/6, hep-ph 9603310
4. J. Ellis, J. L. Lopez, D. V. Nanopoulos, *Supersymmetry and R_b in the light of LEP 1.5*, hep-ph/9512288.
5. D. Garcia, J. Sola, *The Quantum Correlation $R_b - R_c$ in the MSSM: More Hints of Supersymmetry?*, hep-ph/9502317, *Phys. Lett. B***354** (1995) 335,
6. G.L. Kane, R.G. Stuart, J.D. Wells, *A Global Fit of LEP/SLC Data with Light Superpartners*, *Phys. Lett. B***354**(1995) 350, UM-TH-94-16, hep-ph/9505207
7. J.D. Wells, C. Kolda, G.L. Kane, *Implications of $\Gamma(Z \rightarrow b\bar{b})$ for Supersymmetry Searches and Model-Building*, *Phys. Lett. B***338**(1993) 219, UM-TH-94-23, hep-ph/9408228
8. D. Garcia, R. Jimenez, J. Sola, *Supersymmetric Electroweak Renormalization of the Z Width in the MSSM*, *Phys. Lett. B***347** (1995) 309; *Phys. Lett. B***347** (1995) 321
9. D. Garcia, J. Sola, *Matching the low-Energy and the high-Energy Determinations of $\alpha_s(M(Z))$ in the MSSM*, *Phys. Lett. B***357**(1995) 349
10. A. Blondel, these Proceedings, *CERN-internal note LEPEWWG/96-02*, July 1996.
11. CLEO-Collaboration, R. Ammar et al., *Phys. Rev. Lett.* **74**, (1995) 2885
12. W. de Boer, A. Dabelstein, W. Hollik, W. Möhle and U. Schwickerath, *Updated Global Fits of the SM and MSSM to Electroweak Precision Data*, hep-ph/9609209 and references therein.
13. S. Eidelman and F. Jegerlehner, *Z. Phys. C***67** (1995) 585; H. Burkhardt and B. Pietrzyk, *Phys. Lett. B***356**(1995) 398.
14. R.M. Barnett et al., *Phys. Rev. D***54** (1996) 1.
15. N. Watson, for the LEP Coll. at the LEPC-Meeting, CERN, 8 Oct. 1996.
16. M. Schmelling, these Proceedings.
17. W. de Boer, these proceedings and W. de Boer et al., *Combined Fit of Low Energy Constraints to Minimal Supersymmetry and Discovery Potential at LEP II*, hep-ph/9603350 and references therein.
18. W. de Boer et al., *MSSM predictions of the Neutral Higgs Boson Masses And LEP- II Production Cross-Sections*, hep-ph/9603346 and references therein